

IV. AGRICULTURE

4.05. Consumption of Chemical Fertiliser in different Blocks of Cuttack district.

(In 000' M.T.)

Year / Block	Nitrogeous (N)	Phosphatic (P)	Potassic (K)	Total (N + P + K)
1	2	3	4	5
2005-06	9.40	3.02	2.81	15.23
2006-07	10.89	2.71	2.67	16.27
2006-07 BLOCK				
1 Athagarh	0.80	0.15	0.16	1.11
2 Banki	0.73	0.25	0.28	1.26
3 Baramba	0.57	0.13	0.15	0.85
4 Baranga	0.60	0.19	0.17	0.96
5 Cuttack Sadar	0.79	0.21	0.21	1.21
6 Damapara	0.98	0.23	0.27	1.48
7 Kantapada	0.63	0.22	0.18	1.03
8 Mahanga	0.88	0.20	0.22	1.30
9 Narasinghpur	0.95	0.16	0.13	1.24
10 Niali	0.81	0.22	0.18	1.21
11 Nischintakoili	0.90	0.21	0.21	1.32
12 Salipur	1.02	0.20	0.19	1.41
13 Tangi-Choudwar	0.76	0.21	0.17	1.14
14 Tigiria	0.47	0.13	0.15	0.75

Source : Dist. Agriculture Officer, Cuttack.

SOURCE: District Statistics Office, Cuttack Dt:16.03.2010